


## Phenyl derivatives 5

The invention relates to compounds of the formula I




in which

D is absent or

is a saturated, fully or partially unsaturated 3- to 4-membered alkylene chain, in which from 1 to 3 carbon atoms may be replaced by N and/or 1 or 2 carbon atoms may be replaced by 1 or 2 O and/or 1 or 2 S atoms, but where at most up to 3 carbon atoms are replaced and where, in addition, the alkylene chain and/or a nitrogen present therein may be monosubstituted, disubstituted or trisubstituted by Hal, A,  $-\text{[C(R}^3\text{)]}_n\text{-Ar}$ ,  $-\text{[C(R}^3\text{)]}_n\text{-Het}$ ,  $-\text{[C(R}^3\text{)]}_n\text{-cycloalkyl}$ ,  $\text{OR}^2$ ,  $\text{N(R}^2\text{)}_2$ ,  $\text{NO}_2$ ,  $\text{CN}$ ,  $\text{COOR}^2$ ,  $\text{CON(R}^2\text{)}_2$ ,  $\text{NR}^2\text{COA}$ ,  $\text{NR}^2\text{SO}_2\text{A}$ ,  $\text{COR}^2$ ,  $\text{SO}_2\text{NR}^2$  and/or  $\text{S(O)}_m\text{A}$ , and where, furthermore, one  $\text{CH}_2$  group in the alkylene chain may also be replaced by a  $\text{C=O}$  group,

M is a phenyl ring or an aromatic heterocyclic ring, which may contain 1-2 N, O and/or S atoms,

$\text{R}^1$  and  $\text{R}^{1'}$  are each, independently of one another, H, Hal, A,  $\text{OR}^2$ ,  $\text{N(R}^2\text{)}_2$ ,  $\text{NO}_2$ ,  $\text{CN}$ ,  $\text{COOR}^2$ ,  $\text{CON(R}^2\text{)}_2$ ,  $\text{C(=S)N(R}^2\text{)}_2$ ,  $-\text{[C(R}^3\text{)]}_n\text{-Ar}$ ,  $-\text{[C(R}^3\text{)]}_n\text{-Het}$ ,  $-\text{[C(R}^3\text{)]}_n\text{-cycloalkyl}$ ,  $-\text{[C(R}^3\text{)]}_n\text{-N(R}^3\text{)}_2$ ,  $\text{CN}$ ,  $-\text{C(=NH)-NH}_2$  which is unsubstituted or monosubstituted by  $\text{C(=O)R}^3$ ,  $\text{COOR}^3$ ,  $\text{OR}^3$ ,  $\text{OCOR}^3$ ,  $\text{OCOOR}^3$  or by a conventional amino-protecting group, or



- 5  $R^2$  is H, A,  $-[C(R^3)_2]_n\text{-Ar}$ ,  $-[C(R^3)_2]_n\text{-Het}$ ,  $-[C(R^3)_2]_n\text{-cycloalkyl}$ ,  
 $-[C(R^3)_2]_n\text{-N}(R^3)_2$  or  $-[C(R^3)_2]_n\text{-OR}^3$ ,  
 $R^{2'}$  is H, A,  $-[C(R^3)_2]_n\text{-Ar}'$ ,  $-[C(R^3)_2]_n\text{-Het}'$ ,  $-[C(R^3)_2]_n\text{-cycloalkyl}$ ,  
 $-[C(R^3)_2]_n\text{-N}(R^3)_2$  or  $-[C(R^3)_2]_n\text{-OR}^3$ ,  
10  $R^{2''}$  is H, A,  $-[C(R^3)_2]_n\text{-Ar}'$ ,  $-[C(R^3)_2]_n\text{-cycloalkyl}$ ,  $-[C(R^3)_2]_n\text{-N}(R^3)_2$  or  
 $-[C(R^3)_2]_n\text{-OR}^3$ ,  
 $R^3$  is H or A,  
W is a monocyclic or bicyclic saturated, unsaturated or aromatic  
15 carbocyclic or heterocyclic ring having from 1 to 4 N, O and/or S  
atoms, which may be monosubstituted or disubstituted by  $R^2$ ,  
X is  $\text{CONR}^2$ ,  $\text{CONR}^2\text{C}(R^3)_2$ ,  $\text{-C}(R^3)_2\text{NR}^2$ ,  $\text{-C}(R^3)_2\text{NR}^2\text{C}(R^3)_2$ ,  
 $\text{-C}(R^3)_2\text{O-}$ ,  $\text{-C}(R^3)_2\text{OC}(R^3)_2\text{-}$  or  $\text{NR}^2\text{CO}$ ,  
20 Y is alkylene, cycloalkylene, Het-diyl or Ar-diyl,  
T is a monocyclic or bicyclic, saturated, unsaturated or aromatic  
carbocyclic or heterocyclic ring having from 1 to 4 N, O and/or S  
atoms which is monosubstituted or disubstituted by  $=\text{S}$ ,  $=\text{NR}^2$ ,  
 $=\text{N-CN}$ ,  $=\text{N-NO}_2$ ,  $=\text{NOR}^2$ ,  $=\text{NCOR}^2$ ,  $=\text{NCOOR}^2$  or  $=\text{NOCOR}^2$  and  
25 may furthermore be monosubstituted, disubstituted or trisubsti-  
tuted by Hal, A,  $-[C(R^3)_2]_n\text{-Ar}$ ,  $-[C(R^3)_2]_n\text{-Het}$ ,  $-[C(R^3)_2]_n\text{-cycloalkyl}$ ,  
 $\text{OR}^3$ ,  $\text{N}(R^3)_2$ ,  $\text{NO}_2$ ,  $\text{CN}$ ,  $\text{COOR}^2$ ,  $\text{CON}(R^2)_2$ ,  $\text{NR}^2\text{COA}$ ,  
 $\text{NR}^2\text{CON}(R^2)_2$ ,  $\text{NR}^2\text{SO}_2\text{A}$ ,  $\text{COR}^2$ ,  $\text{SO}_2\text{NR}^2$  and/or  $\text{S(O)}_m\text{A}$ ,  
30 A is unbranched or branched alkyl having 1-10 carbon atoms, in  
which one or two  $\text{CH}_2$  groups may be replaced by O or S atoms  
and/or by  $-\text{CH}=\text{CH}-$  groups, and/or in addition 1-7 H atoms may be  
replaced by F,  
35 Ar is phenyl, naphthyl or biphenyl, each of which is unsubstituted or  
monosubstituted, disubstituted or trisubstituted by Hal, A,  $\text{OR}^3$ ,

$N(R^3)_2$ ,  $NO_2$ ,  $CN$ ,  $COOR^3$ ,  $CON(R^3)_2$ ,  $NR^3COA$ ,  $NR^3CON(R^3)_2$ ,  
 $NR^3SO_2A$ ,  $COR^3$ ,  $SO_2N(R^3)_2$ ,  $S(O)_mA$ ,  $-[C(R^3)_2]_n-COOR^{2'}$  or  
 $-O-[C(R^3)_2]_o-COOR^{2'}$ ,

5 Ar' is phenyl or benzyl, each of which is unsubstituted or monosubstituted or disubstituted by Hal,

Het is a monocyclic or bicyclic, saturated, unsaturated or aromatic heterocyclic ring having from 1 to 4 N, O and/or S atoms, which may be unsubstituted or monosubstituted, disubstituted or trisubstituted by carbonyl oxygen,  $=S$ ,  $=N(R^3)_2$ , Hal, A,  $-[C(R^3)_2]_n-Ar$ ,  
10  $-[C(R^3)_2]_n-Het^1$ ,  $-[C(R^3)_2]_n-cycloalkyl$ ,  $-[C(R^3)_2]_n-OR^{2'}$ ,  
 $-[C(R^3)_2]_n-N(R^{2'})_2$ ,  $NO_2$ ,  $CN$ ,  $-[C(R^3)_2]_n-COOR^{2'}$ ,  
 $-[C(R^3)_2]_n-CON(R^{2'})_2$ ,  $-[C(R^3)_2]_n-NR^{2'}COA$ ,  $NR^{2'}CON(R^{2'})_2$ ,  
15  $-[C(R^3)_2]_n-NR^{2'}SO_2A$ ,  $COR^{2'}$ ,  $SO_2NR^{2'}$  and/or  $S(O)_mA$ ,

Het<sup>1</sup> is a monocyclic or bicyclic, saturated, unsaturated or aromatic heterocyclic ring having 1 or 2 N, O and/or S atoms, which may be unsubstituted or monosubstituted or disubstituted by carbonyl oxygen,  $=S$ ,  $=N(R^3)_2$ , Hal, A,  $OR^{2''}$ ,  $N(R^{2''})_2$ ,  $NO_2$ ,  $CN$ ,  $COOR^{2''}$ ,  
20  $CON(R^{2''})_2$ ,  $NR^{2''}COA$ ,  $NR^{2''}CON(R^{2''})_2$ ,  $NR^{2''}SO_2A$ ,  $COR^{2''}$ ,  $SO_2NR^{2''}$  and/or  $S(O)_mA$ ,

Hal is F, Cl, Br or I,

n is 0, 1 or 2,

25 m is 0, 1 or 2,

o is 1, 2 or 3,

and pharmaceutically usable derivatives, solvates and stereoisomers thereof, including mixtures thereof in all ratios.

30 The invention had the object of finding novel compounds having valuable properties, in particular those which can be used for the preparation of medicaments.

35 It has been found that the compounds of the formula I and salts thereof have very valuable pharmacological properties and are well tolerated. In

particular, they exhibit factor Xa-inhibiting properties and can therefore be employed for combating and preventing thromboembolic disorders, such as thrombosis, myocardial infarction, arteriosclerosis, inflammation, apoplexia, angina pectoris, restenosis after angioplasty and claudicatio intermittens.

The compounds of the formula I according to the invention may furthermore be inhibitors of the coagulation factors factor VIIa, factor IXa and thrombin in the blood coagulation cascade.

Aromatic amidine derivatives having an antithrombotic action are disclosed, for example, in EP 0 540 051 B1, WO 98/28269, WO 00/71508, WO 00/71511, WO 00/71493, WO 00/71507, WO 00/71509, WO 00/71512, WO 00/71515 and WO 00/71516. Cyclic guanidines for the treatment of thromboembolic disorders are described, for example, in WO 97/08165. Aromatic heterocyclic compounds having a factor Xa inhibitory activity are disclosed, for example, in WO 96/10022. Substituted N-[(aminoiminomethyl)phenylalkyl]azaheterocyclamides as factor Xa inhibitors are described in WO 96/40679. Pyrazole derivatives are disclosed in WO 01/29006 and WO 02/24690.

The antithrombotic and anticoagulant effect of the compounds according to the invention is attributed to the inhibitory action against activated coagulation protease, known by the name factor Xa, or to the inhibition of other activated serine proteases, such as factor VIIa, factor IXa or thrombin.

Factor Xa is one of the proteases involved in the complex process of blood coagulation. Factor Xa catalyses the conversion of prothrombin into thrombin. Thrombin cleaves fibrinogen into fibrin monomers, which, after crosslinking, make an elementary contribution to thrombus formation. Activation of thrombin may result in the occurrence of thromboembolic dis-

orders. However, inhibition of thrombin may inhibit the fibrin formation involved in thrombus formation.

The inhibition of thrombin can be measured, for example by the method of G. F. Cousins et al. in *Circulation* **1996**, *94*, 1705-1712.

5

Inhibition of factor Xa can thus prevent the formation of thrombin.

The compounds of the formula I according to the invention and their salts engage in the blood coagulation process by inhibiting factor Xa and thus inhibit the formation of thromboses.

10

The inhibition of factor Xa by the compounds according to the invention and the measurement of the anticoagulant and antithrombotic activity can be determined by conventional in-vitro or in-vivo methods. A suitable method is described, for example, by J. Hauptmann et al. in *Thrombosis and Haemostasis* **1990**, *63*, 220-223.

15

The inhibition of factor Xa can be measured, for example by the method of T. Hara et al. in *Thromb. Haemostas.* **1994**, *71*, 314-319.

20

Coagulation factor VIIa initiates the extrinsic part of the coagulation cascade after binding to tissue factor and contributes to the activation of factor X to give factor Xa. Inhibition of factor VIIa thus prevents the formation of factor Xa and thus subsequent thrombin formation.

25

The inhibition of factor VIIa by the compounds according to the invention and the measurement of the anticoagulant and antithrombotic activity can be determined by conventional in-vitro or in-vivo methods. A conventional method for the measurement of the inhibition of factor VIIa is described, for example, by H. F. Ronning et al. in *Thrombosis Research* **1996**, *84*, 73-81.

30

35

Coagulation factor IXa is generated in the intrinsic coagulation cascade and is likewise involved in the activation of factor X to give factor Xa. Inhibition of factor IXa can therefore prevent the formation of factor Xa in a different way.

5 The inhibition of factor IXa by the compounds according to the invention and the measurement of the anticoagulant and antithrombotic activity can be determined by conventional in-vitro or in-vivo methods. A suitable method is described, for example, by J. Chang et al. in *Journal of Biological Chemistry* **1998**, 273, 12089-12094.

The compounds according to the invention may furthermore be used for the treatment of tumours, tumour illnesses and/or tumour metastases.

15 A correlation between tissue factor TF / factor VIIa and the development of various types of cancer has been indicated by T.Taniguchi and N.R. Lemoine in Biomed. Health Res. (2000), 41 (Molecular Pathogenesis of Pancreatic Cancer), 57-59.

20 The publications listed below describe an antitumoural action of TF-VII and factor Xa inhibitors for various types of tumour:

K.M. Donnelly et al. in Thromb. Haemost. 1998; 79: 1041-1047;  
E.G. Fischer et al. in J. Clin. Invest. 104: 1213-1221 (1999);  
B.M. Mueller et al. in J. Clin. Invest. 101: 1372-1378 (1998);  
25 M.E. Bromberg et al. in Thromb. Haemost. 1999; 82: 88-92

The compounds of the formula I can be employed as medicament active ingredients in human and veterinary medicine, in particular for the treatment and prevention of thromboembolic disorders, such as thrombosis, myocardial infarction, arteriosclerosis, inflammation, apoplexia, angina pectoris, restenosis after angioplasty, claudicatio intermittens, venous thrombosis, pulmonary embolism, arterial thrombosis, myocardial ischaemia, unstable angina and strokes based on thrombosis.

30  
35

The compounds according to the invention are also employed for the treatment or prophylaxis of atherosclerotic diseases, such as coronary arterial disease, cerebral arterial disease or peripheral arterial disease.

5 The compounds are also employed in combination with other thrombolytic agents in myocardial infarction, furthermore for prophylaxis for reocclusion after thrombolysis, percutaneous transluminal angioplasty (PTCA) and coronary bypass operations.

10 The compounds according to the invention are furthermore used for the prevention of rethrombosis in microsurgery, furthermore as anticoagulants in connection with artificial organs or in haemodialysis.

The compounds are furthermore used in the cleaning of catheters and medical aids in patients *in vivo*, or as anticoagulants for the preservation of blood, plasma and other blood products *in vitro*. The compounds according to the invention are furthermore used for diseases in which blood coagulation makes a crucial contribution toward the course of the disease or represents a source of secondary pathology, such as, for example, in cancer, including metastasis, inflammatory disorders, including arthritis, and diabetes.

20 The compounds according to the invention are furthermore used for the treatment of migraine (F.Morales-Asin et al., Headache, 40, 2000, 45-47).

25 In the treatment of the disorders described, the compounds according to the invention are also used in combination with other thrombolytically active compounds, such as, for example, with the "tissue plasminogen activator" t-PA, modified t-PA, streptokinase or urokinase. The compounds according to the invention are administered either at the same time as or before or after the other substances mentioned.

30 Particular preference is given to simultaneous administration with aspirin in order to prevent recurrence of the clot formation.

35

The compounds according to the invention are also used in combination with blood platelet glycoprotein receptor (IIb/IIIa) antagonists, which inhibit blood platelet aggregation.


5 The invention relates to the compounds of the formula I and salts thereof and to a process for the preparation of compounds of the formula I according to Claims 1-20 and pharmaceutically usable derivatives, solvates and stereoisomers thereof, characterised in that

10

a) for the preparation of a compound of the formula I in which X is  $\text{CONR}^2$  or  $\text{CONR}^2\text{C}(\text{R}^3)_2$ ,

15

a compound of the formula II



20

in which

L is Cl, Br, I or a free or reactively functionally modified OH group, and R<sup>1</sup>, R<sup>1'</sup>, D, M and W are as defined in Claim 1,

25

with the proviso that any further OH and/or amino group present is protected,

is reacted with a compound of the formula III

30

Z'-Y-T

III

in which

Z' is  $\text{NHR}^2$  or  $\text{NHR}^2\text{C}(\text{R}^3)_2$ ,

35

and R<sup>2</sup>, Y and T are as defined in Claim 1,

and any protecting group is subsequently removed,



b) and/or in that a radical T, R<sup>1</sup> and/or R<sup>1'</sup> in a compound of the formula I is converted into another radical T, R<sup>1</sup> and/or R<sup>1'</sup>

5 by, for example,

- i) converting a sulfanyl compound into an imino compound,
- ii) removing an amino-protecting group,

10

and/or

a base or acid of the formula I is converted into one of its salts.

15

The invention also relates to the optically active forms (stereoisomers), the enantiomers, the racemates, the diastereomers and the hydrates and solvates of these compounds. The term solvates of the compounds is taken to mean adductions of inert solvent molecules onto the compounds which form owing to their mutual attractive force. Solvates are, for example,

20

The term "pharmaceutically usable derivatives" is taken to mean, for example, the salts of the compounds according to the invention and so-called prodrug compounds.

25

The term "prodrug derivatives" is taken to mean compounds of the formula I which have been modified with, for example, alkyl or acyl groups, sugars or oligopeptides and which are rapidly cleaved in the organism to form the active compounds according to the invention.

30

These also include biodegradable polymer derivatives of the compounds according to the invention, as described, for example, in Int. J. Pharm. 115, 61-67 (1995).

35

The invention also relates to mixtures of the compounds of the formula I according to the invention, for example mixtures of two diastereomers, for

example in the ratio 01:01, 01:02, 01:03, 01:04, 01:05, 01:10, 1:100 or 1:1000.

These are particularly preferably mixtures of stereoisomeric compounds.

5 For all radicals which occur more than once, such as, for example, A, their meanings are independent of one another.

Above and below, the radicals and parameters D, M, W, X, Y, T, R<sup>1</sup> and R<sup>1'</sup> are as defined under the formula I, unless expressly stated otherwise.

10

The following abbreviations are used below:

	Ac	acetyl
15	BOC	tert-butoxycarbonyl
	CBZ or Z	benzyloxycarbonyl
	DCCI	dicyclohexylcarbodiimide
	DCM	dichloromethane
	DMF	dimethylformamide
20	EDCI	N-ethyl-N,N'-(dimethylaminopropyl)carbodiimide
	EA	ethyl acetate
	Et	ethyl
	Fmoc	9-fluorenylmethoxycarbonyl
25	HOBt	1-hydroxybenzotriazole
	Me	methyl
	MBHA	4-methylbenzhydrylamine
	Mtr	4-methoxy-2,3,6-trimethylphenylsulfonyl
30	HONSu	N-hydroxysuccinimide
	OBu <sup>t</sup>	tert-butyl ester
	Oct	octanoyl
	OMe	methyl ester
	OEt	ethyl ester
35	POA	phenoxyacetyl
	TFA	trifluoroacetic acid

Trt trityl (triphenylmethyl).

The ring M is preferably phenyl.

5 D, if present, may be monosubstituted, disubstituted or trisubstituted, preferably by Hal, A, OR<sup>2</sup> or N(R<sup>2</sup>)<sub>2</sub>, and/or one CH<sub>2</sub> group in the alkylene chain may also be replaced by a C=O group. Monosubstitution by A or NH<sub>2</sub> is very particularly preferred.

10 D is preferably -CO-NH-CO-, -CO-NH-CH<sub>2</sub>-, -NH-CH=CH-, -O-CH=CH-, -N=CH-O-, -N=CH-NH-, -NH-NH-CO-, -NH-N=N-, -NH-CO-CH<sub>2</sub>-, -NH-CO-O-, -N=CH-S-, -NH-CO-S-, -NH-CO-NH-, -NH-N=CH-, -S-N=CH-,  
15 =C-S-N=, -O-N=CH-, -O-NH-CO-, -NH-O-CO-, -N=CH-CH=CH-, -CH=N-CH=CH-, -N=N-CH=CH-, -N=CH-N=CH-, -N=CH-CH=N-, -N=N-N=CH-, -NH-CO-CH=CH-, -NH-CH=CH-CO-, -NH-CO-CH<sub>2</sub>-CH<sub>2</sub>-, -NH-CH<sub>2</sub>-CH<sub>2</sub>-CO-, -NH-CO-N=CH-, -N=CH-NH-CO-, -NH-CO-NH-CO-, -NH-CO-NH-CH<sub>2</sub>-, -CH=N-N=CH-, -N<sup>-</sup>-S<sup>+</sup>=N-, -O-CH<sub>2</sub>-O-, furthermore  
20 -CH=N-NH-CO-, -CH=CH-NH-, -O-CH<sub>2</sub>CH<sub>2</sub>-O-, -CO-NH-NH-CO-, -N=N-NH-CO-, -O-CO-NH-CH<sub>2</sub>- or -O-CO-NH-CO-, where, in addition, the alkylene chain and/or a nitrogen present therein may be monosubstituted, disubstituted or trisubstituted by A or NH<sub>2</sub>.

25 In a further embodiment, D is preferably a saturated 3- to 4-membered alkylene chain, in which from 1 to 3 carbon atoms may be replaced by N and/or 1 or 2 carbon atoms may be replaced by 1 or 2 O atoms, but where  
30 at most up to 3 carbon atoms are replaced and where, in addition, the alkylene chain and/or or a nitrogen located therein may be monosubstituted or disubstituted by NH<sub>2</sub>, or D is absent.

35 D is particularly preferably -CH=N-CH=CH-, -CH=CH-N=CH-, NH-N=CH-, -CH=N-NH-, -O-N=CH- or -CH=N-O-, where, in addition, D may be monosubstituted by NH<sub>2</sub>, or D is absent.

A is alkyl, is unbranched (linear) or branched, and has 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 carbon atoms. A is preferably methyl, furthermore ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl or tert-butyl, furthermore also pentyl, 1-, 2- or 3-methylbutyl, 1,1-, 1,2- or 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1-, 2-, 3- or 4-methylpentyl, 1,1-, 1,2-, 1,3-, 2,2-, 2,3- or 3,3-dimethylbutyl, 1- or 2-ethylbutyl, 1-ethyl-1-methylpropyl, 1-ethyl-2-methylpropyl, 1,1,2- or 1,2,2-trimethylpropyl, furthermore preferably, for example, trifluoromethyl.

A is very particularly preferably alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms, preferably methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, trifluoromethyl, pentafluoroethyl or 1,1,1-trifluoroethyl.

Alkoxy is preferably, for example, methoxy, ethoxy, propoxy, isopropoxy, butoxy, trifluoromethoxy or cyclopentoxy.

Cycloalkyl is preferably cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl.


Alkylene is preferably methylene, ethylene, propylene, butylene, pentylene or hexylene, furthermore branched alkylene.

COR<sup>2</sup> is for example, CHO or -COA.

-COA (acyl) is preferably acetyl, propionyl, furthermore also butyryl, pentanoyl, hexanoyl or, for example, benzoyl.

Hal is preferably F, Cl or Br, but alternatively I.

R<sup>1</sup> is preferably CN, CONH<sub>2</sub>, CONA<sub>2</sub>, NH<sub>2</sub>, C(=S)NH<sub>2</sub>, CH<sub>2</sub>NH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>, -C(=NH)-NH<sub>2</sub> which is unsubstituted or monosubstituted by OH, OCOA or OCOOA, or



where A is preferably alkyl having 1, 2, 3 or 4 carbon atoms.

R<sup>1</sup> is preferably H, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-N(R<sup>3</sup>)<sub>2</sub>, CON(R<sup>2</sup>)<sub>2</sub>, C(=S)NH<sub>2</sub> or N(R<sup>2</sup>)<sub>2</sub>, particularly preferably H, CH<sub>2</sub>NH<sub>2</sub>, NH<sub>2</sub>, CONH<sub>2</sub> or C(=S)NH<sub>2</sub>.

R<sup>1'</sup> is preferably H.

R<sup>2</sup> is preferably, for example, H or alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms.

R<sup>2'</sup> is preferably, for example, H or alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms.

R<sup>2''</sup> is preferably, for example, H or alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms.

W is a monocyclic or bicyclic saturated, unsaturated or aromatic carbocyclic or heterocyclic ring having from 1 to 4 N, O and/or S atoms, which may be monosubstituted or disubstituted by R<sup>2</sup>.

In a preferred embodiment, W is a monocyclic saturated, unsaturated or aromatic carbocyclic or heterocyclic ring having 1 or 2 N, O and/or S atoms, which may be monosubstituted or disubstituted by R<sup>2</sup>.

In a further preferred embodiment, W is, for example, cyclohexanediyl, cyclopentanediy, phenylene, biphenylene, furandiyl, thiophenediy, pyrrolediyl, imidazolediyl, pyrazolediyl, oxazolediyl, isoxazolediyl, thiazole-diyl, isothiazole-diyl, pyridinediy, pyrimidinediy, pyrrolidinediy, piperidinediy or piperazinediy, each of which is monosubstituted or disubstituted by R<sup>2</sup>.

In a further preferred embodiment, W is, for example, cyclohexanediyl, cyclopentanediy, phenylene, biphenylene, furandiyl, thiophenediy, pyrrolediyl, imidazolediyl, pyrazolediyl, oxazolediyl, isoxazolediyl, thiazole-diyl, isothiazole-diyl, pyridinediy, pyrimidinediy, pyrrolidinediy,

piperidinediyl or piperazinediyl, each of which is unsubstituted or mono-substituted or disubstituted by A or Hal .

W is particularly preferably pyrazolediyl or thiazolediyl, each of which is unsubstituted or monosubstituted by A, CONH<sub>2</sub> or COOA, where A is, in particular, alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms or CF<sub>3</sub>.

X is preferably CONH, CONHCH<sub>2</sub>, CH<sub>2</sub>NH or CH<sub>2</sub>NHCH<sub>2</sub>, very particularly preferably CONH.

Y is preferably alkylene or Ar-diyl, particularly preferably methylene, ethylene, propylene, or 1,4-phenylene which is unsubstituted or monosubstituted by A, Cl or F, furthermore also pyridinediyl, preferably pyridine-2,5-diyl.

Y is, in particular, 1,3- or 1,4-phenylene which is unsubstituted or monosubstituted by methyl, ethyl, propyl, Br, Cl or F, very particularly preferably 1,4-phenylene.

Ar is preferably phenyl, naphthyl or biphenyl, each of which is unsubstituted or monosubstituted, disubstituted or trisubstituted by Hal, A, OH, NH<sub>2</sub>, NO<sub>2</sub>, CN, COOH, CONH<sub>2</sub>, NHCOA, NHCONH<sub>2</sub>, NHSO<sub>2</sub>A, COH, SO<sub>2</sub>NH<sub>2</sub>, S(O)<sub>m</sub>A, -(CH<sub>2</sub>)<sub>n</sub>-COOR<sup>2'</sup> or -O-(CH<sub>2</sub>)<sub>o</sub>-COOR<sup>2'</sup>.

Ar is, for example, unsubstituted phenyl, naphthyl or biphenyl, furthermore preferably phenyl, naphthyl or biphenyl, each of which is monosubstituted, disubstituted or trisubstituted by A, fluorine, chlorine, bromine, iodine, hydroxyl, methoxy, ethoxy, propoxy, butoxy, pentyloxy, hexyloxy, nitro, cyano, formyl, acetyl, propionyl, trifluoromethyl, amino, methylamino, ethylamino, dimethylamino, diethylamino, benzyloxy, sulfonamido, methylsulfonamido, ethylsulfonamido, propylsulfonamido, butylsulfonamido, dimethylsulfonamido, phenylsulfonamido, carboxy, methoxycarbonyl, ethoxycarbonyl or aminocarbonyl.

Het is, for example, 2- or 3-furyl, 2- or 3-thienyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, furthermore preferably 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 3- or 4-pyridazinyl, pyrazinyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 4- or 5-isoindolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5-, 6- or 7-benzisoxazolyl, 2-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 4-, 5-, 6- or 7-benz-2,1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolyl, 1-, 3-, 4-, 5-, 6-, 7- or 8-isoquinolyl, 3-, 4-, 5-, 6-, 7- or 8-quinolyl, 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, 5- or 6-quinoxalyl, 2-, 3-, 5-, 6-, 7- or 8-2H-benzo-1,4-oxazinyl, furthermore preferably 1,3-benzodioxol-5-yl, 1,4-benzodioxan-6-yl, 2,1,3-benzothiadiazol-4- or -5-yl or 2,1,3-benzoxadiazol-5-yl.

The heterocyclic radicals may also be partially or fully hydrogenated. Het can thus also be, for example, 2,3-dihydro-2-, -3-, -4- or -5-furyl, 2,5-dihydro-2-, -3-, -4- or 5-furyl, tetrahydro-2- or -3-furyl, 1,3-dioxolan-4-yl, tetrahydro-2- or -3-thienyl, 2,3-dihydro-1-, -2-, -3-, -4- or -5-pyrrolyl, 2,5-dihydro-1-, -2-, -3-, -4- or -5-pyrrolyl, 1-, 2- or 3-pyrrolidinyl, tetrahydro-1-, -2- or -4-imidazolyl, 2,3-dihydro-1-, -2-, -3-, -4- or -5-pyrazolyl, tetrahydro-1-, -3- or -4-pyrazolyl, 1,4-dihydro-1-, -2-, -3- or -4-pyridyl, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5- or -6-pyridyl, 1-, 2-, 3- or 4-piperidinyl, 2-, 3- or 4-morpholinyl, tetrahydro-2-, -3- or -4-pyranyl, 1,4-dioxanyl, 1,3-dioxan-2-, -4- or -5-yl, hexahydro-1-, -3- or -4-pyridazinyl, hexahydro-1-, -2-, -4- or -5-pyrimidinyl, 1-, 2- or 3-piperazinyl, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5-, -6-, -7- or -8-quinolyl, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5-, -6-, -7- or -8-isoquinolyl, 2-, 3-, 5-, 6-, 7- or 8-3,4-dihydro-2H-benzo-1,4-oxazinyl, furthermore preferably 2,3-methylenedioxyphenyl, 3,4-methylenedioxyphenyl, 2,3-ethylenedioxyphenyl, 3,4-ethylenedioxyphenyl, 3,4-(difluoromethylenedioxy)phenyl, 2,3-dihydrobenzofuran-5- or 6-yl, 2,3-(2-oxo-

methylenedioxy)phenyl or alternatively 3,4-dihydro-2H-1,5-benzodioxepin-6- or -7-yl, furthermore preferably 2,3-dihydrobenzofuranyl or 2,3-dihydro-2-oxofuranyl.

5 Het<sup>1</sup> has the preferred meanings, such as Het.

T is preferably a monocyclic saturated or unsaturated heterocyclic ring having 1 or 2 N and/or O atoms which is monosubstituted or disubstituted  
10 by =S, =NR<sup>2</sup>, =NOR<sup>2</sup>, =N-CN, =N-NO<sub>2</sub>, =NCOR<sup>2</sup>, =NCOOR<sup>2</sup> or =NOCOR<sup>2</sup>, in particular by =S, =NR<sup>2</sup> or =NOR<sup>2</sup>, and may preferably furthermore be monosubstituted or disubstituted by A.

15 T is, in particular, piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl, azepan-1-yl or 2-azabicyclo[2.2.2]octan-2-yl, pyrazol-2-yl, 1,3,4-thiadiazol-3-yl, imidazolidin-1-yl or 1,2-dihydropyrazol-2-yl, each of which is monosubstituted or disubstituted by =NR<sup>2</sup>, =S or =NOR<sup>2</sup> and may preferably furthermore be  
20 monosubstituted or disubstituted by A, CONH<sub>2</sub> or COOA.

T is furthermore particularly preferably, for example, 2-iminopiperidin-1-yl, 2-iminopyrrolidin-1-yl, 2-imino-1*H*-pyridin-1-yl, 3-iminomorpholin-4-yl,  
25 4-imino-1*H*-pyridin-1-yl, 2,6-diiminopiperidin-1-yl, 2-iminopiperazin-1-yl, 2,6-diiminopiperazin-1-yl, 2,5-diiminopyrrolidin-1-yl, 2-imino-1,3-oxazolidin-3-yl, 3-imino-2*H*-pyridazin-2-yl, 2-iminoazepan-1-yl, 2-hydroxy-6-iminopiperazin-1-yl, pyrazol-2-yl, 1,2-dihydropyrazol-2-yl, 2-methoxy-6-iminopiperazin-1-yl, 2-imino-1,3,4-thiadiazol-3-yl or 2-iminoimidazolidin-1-yl,  
30 yl, very particularly preferably 2-iminopiperidin-1-yl, and the corresponding hydroxyimino, alkoxyimino, thioxo and =N-(CH<sub>2</sub>)<sub>1-3</sub>NA'<sub>2</sub> derivatives, where A' is alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms.

35



In a further particularly preferred embodiment, T is, for example, 2-iminopyrrolidin-1-yl, 2-iminopiperidin-1-yl, 2-imino-1,3,4-thiadiazol-3-yl, 2-iminoimidazolidin-1-yl or 3-imino-1,2-dihydropyrazol-2-yl, and the corresponding hydroxyimino, cyanoimino, alkoxyimino and thioxo derivatives, where the heterocyclic radicals may preferably furthermore be monosubstituted or disubstituted by A, CONH<sub>2</sub> or COOA.

In a further particularly preferred embodiment, T is, for example, 2-iminopyrrolidin-1-yl, 2-iminopiperidin-1-yl, 2-thioxopyrrolidin-1-yl, 2-imino-1,3,4-thiadiazol-3-yl, 2-methoxyiminopyrrolidin-1-yl, 2-hydroxyiminopyrrolidin-1-yl, 2-iminoimidazolidin-1-yl or 3-imino-1,2-dihydropyrazol-2-yl, where the heterocyclic radicals may preferably furthermore be monosubstituted or disubstituted by A, CONH<sub>2</sub> or COOA.

The compounds of the formula I may have one or more chiral centres and therefore occur in various stereoisomeric forms. The formula I covers all these forms.

Accordingly, the invention relates in particular to the compounds of the formula I in which at least one of the said radicals has one of the preferred meanings indicated above. Some preferred groups of compounds may be expressed by the following sub-formulae Ia to Iw, which conform to the formula I and in which the radicals not designated in greater detail are as defined under the formula I, but in which

in Ia        D    is absent;

in Ib        M    is a phenyl ring;

in Ic        D    is a saturated, fully or partially unsaturated 3- to 4-membered alkylene chain, in which from 1 to 3 carbon atoms may be replaced by N and/or 1 or 2 carbon atoms

may be replaced by 1 or 2 O and/or 1 or 2 S atoms, but where at most up to 3 carbon atoms are replaced and where, in addition, the alkylene chain and/or a nitrogen present therein may be monosubstituted, disubstituted or trisubstituted by Hal, A, OR<sup>2</sup> or N(R<sup>2</sup>)<sub>2</sub>, and where, furthermore, one CH<sub>2</sub> group in the alkylene chain may also be replaced by a C=O group;

5  
10      in Id      D      is a saturated, fully or partially unsaturated 3- to 4-membered alkylene chain, in which from 1 to 3 carbon atoms may be replaced by N and/or 1 or 2 carbon atoms may be replaced by 1 or 2 O and/or 1 or 2 S atoms, but where at most up to 3 carbon atoms are replaced and where, in addition, the alkylene chain and/or a nitrogen present therein may be monosubstituted, disubstituted or trisubstituted by A or NH<sub>2</sub>;

15  
20      in le      D      is absent or is a saturated 3- to 4-membered alkylene chain, in which from 1 to 3 carbon atoms may be replaced by N and/or 1 or 2 carbon atoms may be replaced by 1 or 2 O atoms, but where at most up to 3 carbon atoms are replaced,  
25  
and where, in addition, the alkylene chain and/or a nitrogen atom located therein may be monosubstituted or disubstituted by NH<sub>2</sub>;

30      in lf      D      is absent or is -CH=N-CH=CH-, -CH=CH-N=CH-, -NH-N=CH-, -CH=N-NH-, -O-N=CH- or -CH=N-O-, and where, in addition, D may be monosubstituted by NH<sub>2</sub>;

35      in lg      R<sup>1</sup>      is H, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-N(R<sup>3</sup>)<sub>2</sub>, CON(R<sup>2</sup>)<sub>2</sub>, C(=S)NH<sub>2</sub> or N(R<sup>2</sup>)<sub>2</sub>,  
R<sup>1'</sup>      is H;

- in lh       $R^1$     is H,  $CH_2NH_2$ ,  $CONH_2$ ,  $C(=S)NH_2$  or  $NH_2$ ,  
               $R^{1'}$     is H;
- 5           in li      W      is a monocyclic saturated, unsaturated or aromatic carbo-  
                              cyclic or heterocyclic ring having 1 or 2 N, O and/or S  
                              atoms, which may be monosubstituted or disubstituted by  
                               $R^2$ ;
- 10           in lj      W      is cyclohexanediyl, cyclopentanediyl, phenylene,  
                              biphenylene, furandiyl, thiophenediyl, pyrrolediyl, imida-  
                              zolediyl, pyrazolediyl, oxazolediyl, isoxazolediyl, thiazole-  
                              diyl, isothiazolediyl, pyridinediyl, pyrimidinediyl, pyrroli-  
                              dinediyl, piperidinediyl or piperazinediyl, each of which is  
                              unsubstituted or monosubstituted or disubstituted by  $R^2$ ;
- 15           in lk      W      is pyrazolediyl, which is unsubstituted or monosubstituted  
                              by A;
- 20           in ll      X      is  $CONH$ ,  $CONHCH_2$ ,  $CH_2NH$  or  $CH_2NHCH_2$ ;
- 25           in lm      X      is  $CONH$ ;
- in ln      Y      is alkylene or Ar-diyl;
- 30           in lo      Y      is phenylene which is unsubstituted or monosubstituted or  
                              disubstituted by A, Br, Cl or F;
- 35           in lp      T      is a monocyclic saturated or unsaturated heterocyclic ring  
                              having from 1 to 3 N, O and/or S atoms, which is mono-  
                              substituted or disubstituted by  $=S$ ,  $=NR^2$ ,  $=NOR^2$ ,  $=N-CN$ ,  
                               $=N-NO_2$ ,  $=NCOR^2$ ,  $=NCOOR^2$  or  $=NOCOR^2$ , and may be

monosubstituted or disubstituted by A,  $\text{CON}(\text{R}^2)_2$  or  $\text{COOR}^2$ ;

- 5            in Iq            T    is a monocyclic saturated or unsaturated heterocyclic ring having from 1 to 3 N, O and/or S atoms, which is monosubstituted or disubstituted by  $=\text{S}$ ,  $=\text{NR}^2$ ,  $=\text{N-CN}$  or  $=\text{NOR}^2$ , and may be monosubstituted or disubstituted by A,  $\text{CON}(\text{R}^2)_2$  or  $\text{COOR}^2$ ;
- 10            in Ir            T    is piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl, azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, pyrazol-2-yl, imidazolidin-1-yl, 1,3,4-thiadiazol-3-yl- or 1,2-dihydropyrazol-2-yl, each of which is monosubstituted or disubstituted by  $=\text{NR}^2$ ,  $=\text{S}$ ,  $=\text{N-CN}$  or  $=\text{NOR}^2$  and which may furthermore be monosubstituted or disubstituted by A,  $\text{CONH}_2$  or  $\text{COOA}$ ,
- 15            in Is            T    is 2-iminopiperidin-1-yl, 2-iminopyrrolidin-1-yl, 2-imino-1*H*-pyridin-1-yl, 3-iminomorpholin-4-yl, 4-imino-1*H*-pyridin-1-yl, 2,6-diiminopiperidin-1-yl, 2-iminopiperazin-1-yl, 2,6-diiminopiperazin-1-yl, 2,5-diiminopyrrolidin-1-yl, 2-imino-1,3-oxazolidin-3-yl, 3-imino-2*H*-pyridazin-2-yl, 2-iminoazepan-1-yl, 2-hydroxy-6-iminopiperazin-1-yl, pyrazol-2-yl, 1,2-dihydropyrazol-2-yl, 2-methoxy-6-iminopiperazin-1-yl, 2-imino-1,3,4-thiadiazol-3-yl, 2-iminoimidazolidin-1-yl, and the corresponding hydroxyimino, alkoxyimino, thioxo and  $=\text{N}-(\text{CH}_2)_{1-3}\text{NA}'_2$  derivatives, where A' is alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms, and where the heterocyclic rings may furthermore be monosubstituted or disubstituted by A,  $\text{CONH}_2$  or  $\text{COOA}$ ;
- 20            in Is            T    is 2-iminopiperidin-1-yl, 2-iminopyrrolidin-1-yl, 2-imino-1*H*-pyridin-1-yl, 3-iminomorpholin-4-yl, 4-imino-1*H*-pyridin-1-yl, 2,6-diiminopiperidin-1-yl, 2-iminopiperazin-1-yl, 2,6-diiminopiperazin-1-yl, 2,5-diiminopyrrolidin-1-yl, 2-imino-1,3-oxazolidin-3-yl, 3-imino-2*H*-pyridazin-2-yl, 2-iminoazepan-1-yl, 2-hydroxy-6-iminopiperazin-1-yl, pyrazol-2-yl, 1,2-dihydropyrazol-2-yl, 2-methoxy-6-iminopiperazin-1-yl, 2-imino-1,3,4-thiadiazol-3-yl, 2-iminoimidazolidin-1-yl, and the corresponding hydroxyimino, alkoxyimino, thioxo and  $=\text{N}-(\text{CH}_2)_{1-3}\text{NA}'_2$  derivatives, where A' is alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms, and where the heterocyclic rings may furthermore be monosubstituted or disubstituted by A,  $\text{CONH}_2$  or  $\text{COOA}$ ;
- 25            in Is            T    is 2-iminopiperidin-1-yl, 2-iminopyrrolidin-1-yl, 2-imino-1*H*-pyridin-1-yl, 3-iminomorpholin-4-yl, 4-imino-1*H*-pyridin-1-yl, 2,6-diiminopiperidin-1-yl, 2-iminopiperazin-1-yl, 2,6-diiminopiperazin-1-yl, 2,5-diiminopyrrolidin-1-yl, 2-imino-1,3-oxazolidin-3-yl, 3-imino-2*H*-pyridazin-2-yl, 2-iminoazepan-1-yl, 2-hydroxy-6-iminopiperazin-1-yl, pyrazol-2-yl, 1,2-dihydropyrazol-2-yl, 2-methoxy-6-iminopiperazin-1-yl, 2-imino-1,3,4-thiadiazol-3-yl, 2-iminoimidazolidin-1-yl, and the corresponding hydroxyimino, alkoxyimino, thioxo and  $=\text{N}-(\text{CH}_2)_{1-3}\text{NA}'_2$  derivatives, where A' is alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms, and where the heterocyclic rings may furthermore be monosubstituted or disubstituted by A,  $\text{CONH}_2$  or  $\text{COOA}$ ;
- 30            in Is            T    is 2-iminopiperidin-1-yl, 2-iminopyrrolidin-1-yl, 2-imino-1*H*-pyridin-1-yl, 3-iminomorpholin-4-yl, 4-imino-1*H*-pyridin-1-yl, 2,6-diiminopiperidin-1-yl, 2-iminopiperazin-1-yl, 2,6-diiminopiperazin-1-yl, 2,5-diiminopyrrolidin-1-yl, 2-imino-1,3-oxazolidin-3-yl, 3-imino-2*H*-pyridazin-2-yl, 2-iminoazepan-1-yl, 2-hydroxy-6-iminopiperazin-1-yl, pyrazol-2-yl, 1,2-dihydropyrazol-2-yl, 2-methoxy-6-iminopiperazin-1-yl, 2-imino-1,3,4-thiadiazol-3-yl, 2-iminoimidazolidin-1-yl, and the corresponding hydroxyimino, alkoxyimino, thioxo and  $=\text{N}-(\text{CH}_2)_{1-3}\text{NA}'_2$  derivatives, where A' is alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms, and where the heterocyclic rings may furthermore be monosubstituted or disubstituted by A,  $\text{CONH}_2$  or  $\text{COOA}$ ;
- 35            in Is            T    is 2-iminopiperidin-1-yl, 2-iminopyrrolidin-1-yl, 2-imino-1*H*-pyridin-1-yl, 3-iminomorpholin-4-yl, 4-imino-1*H*-pyridin-1-yl, 2,6-diiminopiperidin-1-yl, 2-iminopiperazin-1-yl, 2,6-diiminopiperazin-1-yl, 2,5-diiminopyrrolidin-1-yl, 2-imino-1,3-oxazolidin-3-yl, 3-imino-2*H*-pyridazin-2-yl, 2-iminoazepan-1-yl, 2-hydroxy-6-iminopiperazin-1-yl, pyrazol-2-yl, 1,2-dihydropyrazol-2-yl, 2-methoxy-6-iminopiperazin-1-yl, 2-imino-1,3,4-thiadiazol-3-yl, 2-iminoimidazolidin-1-yl, and the corresponding hydroxyimino, alkoxyimino, thioxo and  $=\text{N}-(\text{CH}_2)_{1-3}\text{NA}'_2$  derivatives, where A' is alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms, and where the heterocyclic rings may furthermore be monosubstituted or disubstituted by A,  $\text{CONH}_2$  or  $\text{COOA}$ ;

- in It T is 2-iminopyrrolidin-1-yl, 2-iminopiperidin-1-yl, 2-imino-1,3,4-thiadiazol-3-yl, 2-iminoimidazolidin-1-yl or 3-imino-1,2-dihydropyrazol-2-yl, and the corresponding hydroxyimino, alkoxyimino and thioxo derivatives, where the heterocyclic radicals may furthermore be monosubstituted or disubstituted by A, CONH<sub>2</sub> or COOA;
- 5
- 10 in lu D is absent or is -CH=N-CH=CH-, -CH=CH-N=CH-, -NH-N=CH-, -CH=N-NH-, -O-N=CH- or -CH=N-O-,
- M is a phenyl ring,
- R<sup>1</sup> is H, CH<sub>2</sub>NH<sub>2</sub>, CONH<sub>2</sub>, C(=S)NH<sub>2</sub> or NH<sub>2</sub>,
- 15 R<sup>1'</sup> is H,
- W is a monocyclic saturated, unsaturated or aromatic carbocyclic or heterocyclic ring having 1 or 2 N, O and/or S atoms, which may be monosubstituted or disubstituted by R<sup>2</sup>,
- 20 R<sup>2</sup> is H or alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms,
- R<sup>2'</sup> is H or alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms,
- X is CONH, CONHCH<sub>2</sub>, CH<sub>2</sub>NH or CH<sub>2</sub>NHCH<sub>2</sub>,
- Y is alkylene or Ar-diyl,
- 25 Ar is phenyl, naphthyl or biphenyl, each of which is unsubstituted or monosubstituted, disubstituted or trisubstituted by Hal, A, OH, NH<sub>2</sub>, NO<sub>2</sub>, CN, COOH, CONH<sub>2</sub>, NHCOA, NHCONH<sub>2</sub>, NHSO<sub>2</sub>A, COH, SO<sub>2</sub>NH<sub>2</sub>, S(O)<sub>m</sub>A, -(CH<sub>2</sub>)<sub>n</sub>-COOR<sup>2'</sup> or -O-(CH<sub>2</sub>)<sub>o</sub>-COOR<sup>2'</sup>,
- 30 m and n are each, independently of one another, 0, 1 or 2,
- o is 1, 2 or 3,
- 35 T is piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl, azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, pyrazol-2-yl, 1,3,4-thiadiazol-3-yl, imidazolidin-1-yl or 1,2-dihydro-

pyrazol-2-yl, each of which is monosubstituted or disubstituted by  $=NR^2$ ,  $=N-CN$ ,  $=S$  or  $=NOR^2$  and may furthermore be monosubstituted or disubstituted by A,  $CONH_2$  or  $COOA$ ;

5

- in Iv
- D is absent or is  $-CH=N-CH=CH-$ ,  $-CH=CH-N=CH-$ ,  $-NH-N=CH-$ ,  $-CH=N-NH-$ ,  $-O-N=CH-$  or  $-CH=N-O-$ ,
- M is a phenyl ring,
- 10  $R^1$  is H,  $CH_2NH_2$ ,  $CONH_2$ ,  $C(=S)NH_2$  or  $NH_2$ ,
- $R^{1'}$  is H,
- W is cyclohexanediyl, cyclopentanediy, phenylene, biphenylene, furandiy, thiophenediy, pyrrolediy, imidazolediy, pyrazolediy, oxazolediy, isoxazolediy, thiazolediy, isothiazolediy, pyridinediy, pyrimidinediy, pyrrolidinediy, piperidinediy or piperazinediy, each of which is unsubstituted or monosubstituted or disubstituted by  $R^2$ ,
- 15  $R^2$  is H or alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms,
- 20  $R^{2'}$  is H or alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms,
- X is  $CONH$ ,  $CONHCH_2$ ,  $CH_2NH$  or  $CH_2NHCH_2$ ,
- Y is phenylene which is unsubstituted or monosubstituted or disubstituted by A, Br, Cl or F,
- 25 A is unbranched or branched alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms and/or in addition 1-7 H atoms may be replaced by F,
- 30 T is piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl, azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, pyrazol-2-yl, 1,3,4-thiadiazol-3-yl, imidazolidin-1-yl or 1,2-dihydropyrazol-2-yl, each of which is monosubstituted or disubstituted by  $=NR^2$ ,  $=N-CN$ ,  $=S$  or  $=NOR^2$  and may furthermore be monosubstituted or disubstituted by A,  $CONH_2$  or  $COOA$ ;
- 35

in lw

5. D is absent or is  $-\text{CH}=\text{N}-\text{CH}=\text{CH}-$ ,  $-\text{CH}=\text{CH}-\text{N}=\text{CH}-$ ,  $-\text{NH}-\text{N}=\text{CH}-$ ,  $-\text{CH}=\text{N}-\text{NH}-$ ,  $-\text{O}-\text{N}=\text{CH}-$  or  $-\text{CH}=\text{N}-\text{O}-$ ,  
M is a phenyl ring,  
 $\text{R}^1$  is H,  $\text{CH}_2\text{NH}_2$ ,  $\text{CONH}_2$ ,  $\text{C}(=\text{S})\text{NH}_2$  or  $\text{NH}_2$ ,  
 $\text{R}^{1'}$  is H,  
W is pyrazolediyl or thiazolediyl, each of which is unsubstituted or monosubstituted by A,  
10 X is  $\text{CONH}$ ,  
Y is phenylene which is unsubstituted or monosubstituted or disubstituted by A, Br, Cl or F,  
T is 2-iminopyrrolidin-1-yl, 2-iminopiperidin-1-yl, 2-imino-1,3,4-thiadiazol-3-yl, 2-iminoimidazolidin-1-yl or 3-imino-1,2-dihydropyrazol-2-yl, and the corresponding hydroxyimino, cyanoimino, alkoxyimino and thioxo derivatives, where the heterocyclic radicals may furthermore be monosubstituted or disubstituted by A,  $\text{CONH}_2$  or  $\text{COOA}$ ,  
15 A is unbranched or branched alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms and/or in addition 1-7 H atoms may be replaced by F;  
20

25 and pharmaceutically usable derivatives, solvates and stereoisomers thereof, including mixtures thereof in all ratios.


Furthermore, the invention relates, in particular, to the compounds of the formula I in which at least one of the said radicals has one of the preferred meanings indicated above. Some preferred groups of compounds may be expressed by the following sub-formulae Ia<sub>a</sub> to Ia<sub>c</sub>, which conform to the formula I and in which the radicals not designated in greater detail are as defined under the formula I, but in which

in laa      D      is absent.

M is phenyl,

R<sup>1</sup> is -C(=NH)-NH<sub>2</sub> which is unsubstituted or monosubstituted by OH, or

5



R<sup>1'</sup> is H,

10

W is a monocyclic saturated, unsaturated or aromatic carbocyclic or heterocyclic ring having 1 or 2 N, O and/or S atoms, which may be monosubstituted or disubstituted by R<sup>2</sup>,

15

R<sup>2</sup> is H or alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms,

R<sup>2'</sup> is H or alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms,

X is CONH, CONHCH<sub>2</sub>, CH<sub>2</sub>NH or CH<sub>2</sub>NHCH<sub>2</sub>,

Y is alkylene or Ar-diyl,

20

Ar is phenyl, naphthyl or biphenyl, each of which is unsubstituted or monosubstituted, disubstituted or trisubstituted by Hal, A, OH, NH<sub>2</sub>, NO<sub>2</sub>, CN, COOH, CONH<sub>2</sub>, NHCOA, NHCONH<sub>2</sub>, NHSO<sub>2</sub>A, COH, SO<sub>2</sub>NH<sub>2</sub>, S(O)<sub>m</sub>A, -(CH<sub>2</sub>)<sub>n</sub>-COOR<sup>2'</sup> or -O-(CH<sub>2</sub>)<sub>o</sub>-COOR<sup>2'</sup>,

25

m and n are each, independently of one another, 0, 1 or 2,

o is 1, 2 or 3,

30

T is piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl, azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, pyrazol-2-yl or 1,2-dihydropyrazol-2-yl, each of which is monosubstituted or disubstituted by =NR<sup>2</sup>, =S or =NOR<sup>2</sup> and may furthermore be monosubstituted or disubstituted by A;

35


in lab

D is absent,

M is phenyl,



$R^1$  is  $-C(=NH)-NH_2$  which is unsubstituted or monosubstituted by OH, or



$R^{1'}$  is H,

W is cyclohexanediyl, cyclopentanediy, phenylene, biphenylene, furandiyl, thiophenediyl, pyrrolediyl, imidazolediyl, pyrazolediyl, oxazolediyl, isoxazolediyl, thiazole-diyl, isothiazole-diyl, pyridinediyl, pyrimidinediyl, pyrrolidinediyl, piperidinediyl or piperazinediyl, each of which is unsubstituted or monosubstituted or disubstituted by  $R^2$ ,

$R^2$  is H or alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms,

$R^2$  is H or alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms,

X is CONH, CONHCH<sub>2</sub>, CH<sub>2</sub>NH or CH<sub>2</sub>NHCH<sub>2</sub>,

Y is phenylene which is unsubstituted or monosubstituted or disubstituted by A, Br, Cl or F,


A is unbranched or branched alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms and/or in addition 1-7 H atoms may be replaced by F,

T is piperidin-1-yl, pyrrolidin-1-yl, 1H-pyridin-1-yl, morpholin-4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2H-pyridazin-2-yl, azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, pyrazol-2-yl or 1,2-dihydropyrazol-2-yl, each of which is monosubstituted or disubstituted by  $=NR^2$ ,  $=S$  or  $=NOR^2$  and may furthermore be monosubstituted or disubstituted by A;

in lac D is absent,

M is phenyl,

$R^1$  is CN, NH<sub>2</sub>, CH<sub>2</sub>NH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>,  $-C(=NH)-NH_2$  which is unsubstituted or monosubstituted by OH,



5 R<sup>1'</sup> is H,  
W is pyrazolediyl which is unsubstituted or monosubstituted  
by A,  
X is CONH,  
10 Y is phenylene which is unsubstituted or monosubstituted or  
disubstituted by A, Br, Cl or F,  
T is 2-iminopyrrolidin-1-yl, 2-iminopiperidin-1-yl, 2-imino-  
1,3,4-thiadiazol-3-yl or 3-imino-1,2-dihydropyrazol-2-yl, and  
the corresponding hydroxyimino, alkoxyimino and thioxo  
15 derivatives, where the heterocyclic radicals may  
furthermore be monosubstituted or disubstituted by A,  
A is unbranched or branched alkyl having 1, 2, 3, 4, 5 or 6  
carbon atoms and/or in addition 1-7 H atoms may be  
20 replaced by F;

and pharmaceutically usable derivatives, solvates and stereoisomers thereof, including mixtures thereof in all ratios.


25 The compounds of the formula I and also the starting materials for the  
preparation are, in addition, prepared by methods known per se, as  
described in the literature (for example in the standard works, such as  
Houben-Weyl, Methoden der organischen Chemie [Methods of Organic  
30 Chemistry], Georg-Thieme-Verlag, Stuttgart), to be precise under reaction  
conditions which are known and suitable for the said reactions. Use can  
also be made here of variants which are known per se, but are not men-  
tioned here in greater detail.

35


If desired, the starting materials can also be formed in situ so that they are not isolated from the reaction mixture, but instead are immediately converted further into the compounds of the formula I.

- 5 The starting compounds of the formulae II and III are generally known. If they are novel, they can, however, be prepared by methods known per se.

10 **All** compounds of the following formula VI (where R = H or methyl; n = 3, 4 or 5) can be synthesised in accordance with the following scheme:




- 20 For example, synthesis of 1-(4-amino-2-methylphenyl)piperidine-2-thione:




Alternative synthesis:

- 28 -




5




10

Synthesis of the phenylpiperidinethione unit without a methyl group:



15



20

Compounds of the formula I can preferably be obtained by reacting compounds of the formula II with compounds of the formula III.

25

The reaction is generally carried out in an inert solvent.

30

Examples of suitable inert solvents are hydrocarbons, such as hexane, petroleum ether, benzene, toluene or xylene; chlorinated hydrocarbons, such as trichloroethylene, 1,2-dichloroethane, tetrachloromethane, chloroform or dichloromethane; alcohols, such as methanol, ethanol, isopropanol, n-propanol, n-butanol or tert-butanol; ethers, such as diethyl ether, diisopropyl ether, tetrahydrofuran (THF) or dioxane; glycol ethers, such as ethylene glycol monomethyl or monoethyl ether or ethylene glycol dimethyl ether (diglyme); ketones, such as acetone or butanone; amides, such as acetamide, dimethylacetamide or dimethylformamide (DMF);

35

nitriles, such as acetonitrile; sulfoxides, such as dimethyl sulfoxide (DMSO); carbon disulfide; carboxylic acids, such as formic acid or acetic acid; nitro compounds, such as nitromethane or nitrobenzene; esters, such as ethyl acetate, or mixtures of the said solvents.

5

In the compounds of the formula II, L is preferably Cl, Br, I or a free or reactively modified OH group, such as, for example, an activated ester, an imidazolidine or alkylsulfonyloxy having 1-6 carbon atoms (preferably methylsulfonyloxy or trifluoromethylsulfonyloxy) or arylsulfonyloxy having 6-10 carbon atoms (preferably phenyl- or p-tolylsulfonyloxy).

10

Radicals of this type for activation of the carboxyl group in typical acylation reactions are described in the literature (for example in the standard works, such as Houben-Weyl, Methoden der organischen Chemie [Methods of Organic Chemistry], Georg-Thieme-Verlag, Stuttgart).

15

Activated esters are advantageously formed in situ, for example through addition of HOBt or N-hydroxysuccinimide.

20

The reaction is generally carried out in an inert solvent, in the presence of an acid-binding agent, preferably an organic base, such as DIPEA, triethylamine, dimethylaniline, pyridine, N-methylmorpholine or quinoline, or an excess of the carboxyl component of the formula II.

25

It may also be favourable to add an alkali or alkaline earth metal hydroxide, carbonate or bicarbonate or another salt of a weak acid of the alkali or alkaline earth metals, preferably of potassium, sodium, calcium or caesium.

30

Depending on the conditions used, the reaction time is between a few minutes and 14 days, and the reaction temperature is between about -30° and 140°, normally between -10° and 90°, in particular between about 0° and about 70°.

Suitable inert solvents are those mentioned above.

35

Compounds of the formula I can furthermore be obtained by liberating compounds of the formula I from one of their functional derivatives by treatment with a solvolysing or hydrogenolysing agent.

5 Preferred starting materials for the solvolysis or hydrogenolysis are those which conform to the formula I, but contain corresponding protected amino and/or hydroxyl groups instead of one or more free amino and/or hydroxyl groups, preferably those which carry an amino-protecting group instead of  
10 an H atom bonded to an N atom, in particular those which carry an R'-N group, in which R' is an amino-protecting group, instead of an HN group, and/or those which carry an hydroxyl-protecting group instead of the H atom of an hydroxyl group, for example those which conform to the  
15 formula I, but carry a -COOR" group, in which R" is an hydroxyl-protecting group, instead of a -COOH group.

Preferred starting materials are also the oxadiazole derivatives, which can be converted into the amidino compounds.

20 It is also possible for a plurality of – identical or different – protected amino and/or hydroxyl groups to be present in the molecule of the starting material. If the protecting groups present are different from one another, they can in many cases be cleaved off selectively.

25 The term "amino-protecting group" is known in general terms and relates to groups which are suitable for protecting (blocking) an amino group against chemical reactions, but which are easy to remove after the desired  
30 chemical reaction has been carried out elsewhere in the molecule. Typical of such groups are, in particular, unsubstituted or substituted acyl, aryl, aralkoxymethyl or aralkyl groups. Since the amino-protecting groups are removed after the desired reaction (or reaction sequence), their type and size is furthermore not crucial; however, preference is given to those hav-  
35 ing 1-20, in particular 1-8, carbon atoms. The term "acyl group" is to be understood in the broadest sense in connection with the present process.

It includes acyl groups derived from aliphatic, araliphatic, aromatic or heterocyclic carboxylic acids or sulfonic acids, and, in particular, alkoxy-carbonyl, aryloxy-carbonyl and especially aralkoxycarbonyl groups. Examples of such acyl groups are alkanoyl, such as acetyl, propionyl and butyryl; aralkanoyl, such as phenylacetyl; aroyl, such as benzoyl and tolyl; aryloxyalkanoyl, such as POA; alkoxy-carbonyl, such as methoxycarbonyl, ethoxycarbonyl, 2,2,2-trichloroethoxycarbonyl, BOC (tert-butoxycarbonyl) and 2-iodoethoxycarbonyl; aralkoxycarbonyl, such as CBZ ("carbo-benzoyl"), 4-methoxybenzyloxycarbonyl and FMOC; and arylsulfonyl, such as Mtr. Preferred amino-protecting groups are BOC and Mtr, furthermore CBZ, Fmoc, benzyl and acetyl.

The term "hydroxyl-protecting group" is likewise known in general terms and relates to groups which are suitable for protecting a hydroxyl group against chemical reactions, but are easily removable after the desired chemical reaction has been carried out elsewhere in the molecule. Typical of such groups are the above-mentioned unsubstituted or substituted aryl, aralkyl or acyl groups, furthermore also alkyl groups. The nature and size of the hydroxyl-protecting groups are not crucial since they are removed again after the desired chemical reaction or reaction sequence; preference is given to groups having 1-20, in particular 1-10, carbon atoms. Examples of hydroxyl-protecting groups are, inter alia, benzyl, 4-methoxybenzyl, p-nitrobenzoyl, p-toluenesulfonyl, tert-butyl and acetyl, where benzyl and tert-butyl are particularly preferred.

The compounds of the formula I are liberated from their functional derivatives – depending on the protecting group used – for example using strong acids, advantageously using TFA or perchloric acid, but also using other strong inorganic acids, such as hydrochloric acid or sulfuric acid, strong organic carboxylic acids, such as trichloroacetic acid, or sulfonic acids, such as benzene- or p-toluenesulfonic acid. The presence of an additional inert solvent is possible, but is not always necessary. Suitable inert sol-

vents are preferably organic, for example carboxylic acids, such as acetic acid, ethers, such as tetrahydrofuran or dioxane, amides, such as DMF, halogenated hydrocarbons, such as dichloromethane, furthermore also alcohols, such as methanol, ethanol or isopropanol, and water. Mixtures of the above-mentioned solvents are furthermore suitable. TFA is preferably used in excess without addition of a further solvent, and perchloric acid is preferably used in the form of a mixture of acetic acid and 70% perchloric acid in the ratio 9:1. The reaction temperatures for the cleavage are advantageously between about 0 and about 50°, preferably between 15 and 30° (room temperature).

The BOC, OBut and Mtr groups can, for example, preferably be cleaved off using TFA in dichloromethane or using approximately 3 to 5N HCl in dioxane at 15-30°, and the FMOC group can be cleaved off using an approximately 5 to 50% solution of dimethylamine, diethylamine or piperidine in DMF at 15-30°.

Protecting groups which can be removed hydrogenolytically (for example CBZ, benzyl or the liberation of the amidino group from its oxadiazole derivative) can be cleaved off, for example, by treatment with hydrogen in the presence of a catalyst (for example a noble-metal catalyst, such as palladium, advantageously on a support, such as carbon). Suitable solvents here are those indicated above, in particular, for example, alcohols, such as methanol or ethanol, or amides, such as DMF. The hydrogenolysis is generally carried out at temperatures between about 0 and 100° and pressures between about 1 and 200 bar, preferably at 20-30° and 1-10 bar. Hydrogenolysis of the CBZ group succeeds well, for example, on 5 to 10% Pd/C in methanol or using ammonium formate (instead of hydrogen) on Pd/C in methanol/DMF at 20-30°.

Examples of suitable inert solvents are hydrocarbons, such as hexane, petroleum ether, benzene, toluene or xylene; chlorinated hydrocarbons,



such as trichloroethylene, 1,2-dichloroethane, tetrachloromethane, trifluoromethylbenzene, chloroform or dichloromethane; alcohols, such as methanol, ethanol, isopropanol, n-propanol, n-butanol or tert-butanol; ethers, such as diethyl ether, diisopropyl ether, tetrahydrofuran (THF) or dioxane; glycol ethers, such as ethylene glycol monomethyl or monoethyl ether or ethylene glycol dimethyl ether (diglyme); ketones, such as acetone or butanone; amides, such as acetamide, dimethylacetamide, N-methylpyrrolidone (NMP) or dimethylformamide (DMF); nitriles, such as acetonitrile; sulfoxides, such as dimethyl sulfoxide (DMSO); carbon disulfide; carboxylic acids, such as formic acid or acetic acid; nitro compounds, such as nitromethane or nitrobenzene; esters, such as ethyl acetate, or mixtures of the said solvents.

Esters can be saponified, for example, using acetic acid or using NaOH or KOH in water, water/THF or water/dioxane, at temperatures between 0 and 100°.

Free amino groups can furthermore be acylated in a conventional manner using an acid chloride or anhydride or alkylated using an unsubstituted or substituted alkyl halide or reacted with  $\text{CH}_3\text{-C(=NH)-OEt}$ , advantageously in an inert solvent, such as dichloromethane or THF and/or in the presence of a base, such as triethylamine or pyridine, at temperatures between -60 and +30°.

A base of the formula I can be converted into the associated acid-addition salt using an acid, for example by reaction of equivalent amounts of the base and the acid in an inert solvent, such as ethanol, followed by evaporation. Suitable acids for this reaction are, in particular, those which give physiologically acceptable salts. Thus, it is possible to use inorganic acids, for example sulfuric acid, nitric acid, hydrohalic acids, such as hydrochloric acid or hydrobromic acid, phosphoric acids, such as orthophosphoric acid, or sulfamic acid, furthermore organic acids, in particular aliphatic, alicyclic,

5 araliphatic, aromatic or heterocyclic monobasic or polybasic carboxylic, sulfonic or sulfuric acids, for example formic acid, acetic acid, propionic acid, pivalic acid, diethylacetic acid, malonic acid, succinic acid, pimelic acid, fumaric acid, maleic acid, lactic acid, tartaric acid, malic acid, citric acid, gluconic acid, ascorbic acid, nicotinic acid, isonicotinic acid, methane- or ethanesulfonic acid, ethanedisulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, naphthalenemono- and -disulfonic acids, and laurylsulfuric acid. Salts with physiologically  
10 unacceptable acids, for example picrates, can be used for the isolation and/or purification of the compounds of the formula I.

15 On the other hand, compounds of the formula I can be converted into the corresponding metal salts, in particular alkali metal or alkaline earth metal salts, or into the corresponding ammonium salts using bases (for example sodium hydroxide, potassium hydroxide, sodium carbonate or potassium carbonate). It is also possible to use physiologically acceptable organic bases, such as, for example, ethanolamine.

20 Compounds of the formula I according to the invention may be chiral owing to their molecular structure and may accordingly occur in various enantiomeric forms. They can therefore exist in racemic or in optically active form.

25 Since the pharmaceutical activity of the racemates or stereoisomers of the compounds according to the invention may differ, it may be desirable to use the enantiomers. In these cases, the end product or even the intermediates can be separated into enantiomeric compounds by chemical or  
30 physical measures known to the person skilled in the art or even employed as such in the synthesis.

35 In the case of racemic amines, diastereomers are formed from the mixture by reaction with an optically active resolving agent. Examples of suitable resolving agents are optically active acids, such as the R and S forms of

5 tartaric acid, diacetyltartaric acid, dibenzoyltartaric acid, mandelic acid, malic acid, lactic acid, suitable N-protected amino acids (for example N-benzoylproline) or N-benzenesulfonylproline), or the various optically active camphorsulfonic acids. Also advantage is chromatographic enantio-  
mer resolution with the aid of an optically active resolving agent (for example dinitrobenzoylphenylglycine, cellulose triacetate or other deriva-  
tives of carbohydrates or chirally derivatised methacrylate polymers immo-  
bilised on silica gel). Examples of suitable eluents for this purpose are  
10 aqueous or alcoholic solvent mixtures, such as, for example, hexane/ iso-  
propanol/acetonitrile, for example in the ratio 82:15:3.

15 The invention furthermore relates to the use of compounds of the formula I and/or their physiologically acceptable salts for the preparation of a medi-  
cament (pharmaceutical preparation), in particular by non-chemical  
methods. They can be converted here into a suitable dosage form together  
with at least one solid, liquid and/or semiliquid excipient or assistant and, if  
desired, in combination with one or more further active ingredients.

20 The invention furthermore relates to medicaments comprising at least one  
compound of the formula I and/or its pharmaceutically usable derivatives,  
solvates and stereoisomers, including mixtures thereof in all ratios, and  
25 optionally excipients and/or assistants.

30 These preparations can be used as medicaments in human or veterinary  
medicine. Suitable excipients are organic or inorganic substances which  
are suitable for enteral (for example oral), parenteral or topical administra-  
tion and do not react with the novel compounds, for example water, vege-  
table oils, benzyl alcohols, alkylene glycols, polyethylene glycols, glycerol  
triacetate, gelatin, carbohydrates, such as lactose or starch, magnesium  
stearate, talc or vaseline. Suitable for oral administration are, in particular,  
35 tablets, pills, coated tablets, capsules, powders, granules, syrups, juices or  
drops, suitable for rectal administration are suppositories, suitable for par-

enteral administration are solutions, preferably oil-based or aqueous solutions, furthermore suspensions, emulsions or implants, and suitable for topical application are ointments, creams or powders or also as nasal sprays. The novel compounds may also be lyophilised and the resultant lyophilisates used, for example, to prepare injection preparations. The preparations indicated may be sterilised and/or comprise assistants, such as lubricants, preservatives, stabilisers and/or wetting agents, emulsifying agents, salts for modifying the osmotic pressure, buffer substances, colorants and flavours and/or a plurality of further active ingredients, for example one or more vitamins.

The compounds of the formula I and their physiologically acceptable salts can be used for combating thromboembolic diseases, such as thrombosis, myocardial infarction, arteriosclerosis, inflammation, apoplexia, angina pectoris, restenosis after angioplasty, claudicatio intermittens, tumours, tumour diseases and/or tumour metastases.

In general, the substances according to the invention are preferably administered in doses between about 1 and 500 mg, in particular between 5 and 100 mg, per dosage unit. The daily dose is preferably between about 0.02 and 10 mg/kg of body weight. However, the specific dose for each patient depends on a wide variety of factors, for example on the efficacy of the specific compound employed, on the age, body weight, general state of health, sex, on the diet, on the time and method of administration, on the excretion rate, medicament combination and severity of the particular disease to which the therapy applies. Oral administration is preferred.

The invention furthermore relates to medicaments comprising at least one compound of the formula I and/or its pharmaceutically usable derivatives, solvates and stereoisomers, including mixtures thereof in all ratios, and at least one further medicament active ingredient.

The invention also relates to a set (kit) consisting of separate packs of

(a) an effective amount of a compound of the formula I and/or its  
pharmaceutically usable derivatives, solvates and stereoisomers,  
including mixtures thereof in all ratios,  
and

(b) an effective amount of a further medicament active ingredient.

The set comprises suitable containers, such as boxes, individual bottles,  
bags or ampoules. The set may, for example, comprise separate  
ampoules each containing an effective amount of a compound of the  
formula I and/or its pharmaceutically usable derivatives, solvates and  
stereoisomers, including mixtures thereof in all ratios,  
and an effective amount of a further medicament active ingredient in dis-  
solved or lyophilised form.

The invention furthermore relates to the use of compounds of the formula I  
and/or their pharmaceutically usable derivatives, solvates and stereo-  
isomers, including mixtures thereof in all ratios,  
for the preparation of a medicament for the treatment of thromboses, myo-  
cardial infarction, arteriosclerosis, inflammation, apoplexia, angina pecto-  
ris, restenosis after angioplasty, claudicatio intermittens, migraine,  
tumours, tumour diseases and/or tumour metastases,  
in combination with at least one further medicament active ingredient.


Above and below, all temperatures are given in °C. In the following exam-  
ples, 'conventional work-up' means that water is added if necessary, the  
pH is adjusted, if necessary, to between 2 and 10, depending on the con-  
stitution of the end product, the mixture is extracted with ethyl acetate or  
dichloromethane, the phases are separated, the organic phase is dried  
over sodium sulfate and evaporated, and the product is purified by chro-

matography on silica gel and/or by crystallisation. R<sub>f</sub> values on silica gel;  
eluent: ethyl acetate/methanol 9:1.

Mass spectrometry (MS): EI (electron impact ionisation) M<sup>+</sup>  
FAB (fast atom bombardment) (M+H)<sup>+</sup>  
ESI (electrospray ionisation) (M+H)<sup>+</sup> (unless  
stated otherwise)

### Example 1


*Preparation of an amine unit:*



10 g (48.95 mmol) of 1-(4-amino-2-methylphenyl)piperidin-2-one are  
heated to the boil in 70 ml of anhydrous toluene together with 9.9 g  
(24.48 mmol) of 2,4-bis(4-methoxyphenyl)-1,3,2,4-dithiadiphosphetane  
2,4-disulfide (Lawesson's reagent). After 40 minutes, the solvent is  
removed, and the residue is taken up in dichloromethane (DCM)/1 M  
aqueous hydrochloric acid. After repeated washing with DCM, a pH of 12  
is set using conc. sodium hydroxide solution. Extraction with DCM, drying  
over Na<sub>2</sub>SO<sub>4</sub> and evaporation of the solvent give 9.25 g (41.98 mmol) of  
1-(4-amino-2-methylphenyl)piperidine-2-thione.

### Example 2

*Preparation of an amine unit:*



5

10

2.1 15 g (78.8 mmol) of 1-(4-aminophenyl)piperidin-2-one are heated to the boil together with 16.0 g (39.5 mmol) of 2,4-bis(4-methoxyphenyl)-1,3,2,4-dithiadiphosphetane 2,4-disulfide (Lawesson's reagent) in 100 ml of anhydrous toluene. After 45 minutes, the solvent is evaporated, and the residue is taken up in dichloromethane and 2 N HCl. The aqueous phase is extracted three times with dichloromethane and adjusted to a pH of 12 using conc. NaOH. Extraction with dichloromethane, drying over sodium sulfate and evaporation of the solvent give 1-(4-aminophenyl)piperidine-2-thione as a colourless solid, ESI 207.

20

25

2.2 1.25 ml (20.0 mmol) of iodomethane are added to a solution of 3.74 g (18.1 mmol) of 1-(4-aminophenyl)piperidine-2-thione in 30 ml of acetone, and the mixture is stirred at room temperature for 48 hours. The reaction mixture is evaporated, giving 1-(4-aminophenyl)-6-methylsulfanylium 2,3,4,5-tetrahydropyridinium iodide as a brownish solid; ESI 221.

30

2.3 3.5 ml (25 mmol) of triethylamine are added to a solution of 2.68 g (12.1 mmol) of 1-(4-aminophenyl)-6-methylsulfanylium 2,3,4,5-tetrahydropyridinium iodide and 1.01 g (12.1 mmol) of O-methylhydroxylammonium chloride in 30 ml of ethanol, and the mixture is stirred at room temperature for 20 hours. The reaction mixture is evaporated and taken up in water, and the resultant precipitate is filtered off, giving 1-(4-aminophenyl)piperidin-2-one O-methyl oxime as a colourless solid; ESI 220.

35


**Example 3**

*Preparation of N-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-(3-aminomethyl-phenyl)-5-trifluoromethyl-2H-pyrazole-3-carboxamide ("AA")*

5

10

15



20 3.1 430 mg (1.53 mmol) of 2-(3-cyanophenyl)-5-trifluoromethyl-2H-pyrazole-3-carboxylic acid are hydrogenated on Raney nickel in 10 ml of 10% ammonia in methanol. Work-up and precipitation using ether give 370 mg of 2-(3-aminomethylphenyl)-5-trifluoromethyl-2H-pyrazole-3-carboxylic acid.

25 3.2 370 mg (1.3 mmol) of 2-(3-aminomethylphenyl)-5-trifluoromethyl-2H-pyrazole-3-carboxylic acid are suspended in 6 ml of water, and 303 mg (2.86 mmol) of sodium carbonate are added. A solution of 312 mg (1.43 mmol) of di-*tert*-butyl dicarbonate in 12 ml of 1,4-dioxane is subsequently added dropwise to the reaction mixture, and the mixture is stirred at room temperature for 18 hours. Conventional work-up gives 360 mg of 2-[3-(*tert*-butoxycarbonylaminomethyl)phenyl]-5-trifluoromethyl-2H-pyrazole-3-carboxylic acid as product.

30 3.3 360 mg (0.93 mmol) of 2-[3-(*tert*-butoxycarbonylaminomethyl)phenyl]-5-trifluoromethyl-2H-pyrazole-3-carboxylic acid in 2.5 ml of N,N-



dimethylformamide are reacted with 211 mg (1.0 mmol) of 4-(2-imino-  
pyrrolidin-1-yl)phenylamine hydrochloride with addition of 192 mg (1 mmol)  
of N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride and  
153 mg (1.0 mmol) of 1-hydroxybenzotriazole hydrate. Work-up gives  
5 333 mg of *tert*-butyl (3-{5-[4-(2-iminopyrrolidin-1-yl)phenylcarbamoyl]-3-  
trifluoromethylpyrazol-1-yl}benzyl)carbamate.

3.4 333 mg (0.61 mmol) of *tert*-butyl (3-{5-[4-(2-iminopyrrolidin-1-  
yl)phenylcarbamoyl]-3-trifluoromethylpyrazol-1-yl}benzyl)carbamate are  
10 dissolved in 1 ml of ethanol, and 5 ml of HCl in ether are added. Evapora-  
tion and precipitation using ether give 289 mg of N-[4-(2-iminopyrrolidin-1-  
yl)phenyl]-2-(3-aminomethylphenyl)-5-trifluoromethyl-2*H*-pyrazole-3-  
carboxamide ("AA") in the form of the dihydrochloride as product.

#### 15 Example 4

4.1 Analogously to Example 3, reaction of  
2-(3-aminocarbonylphenyl)-5-trifluoromethyl-2*H*-pyrazole-3-  
20 carboxylic acid with  
1-(4-aminophenyl)pyrrolidine-2-thione gives the compound  
N-[4-(2-thioxopyrrolidin-1-yl)phenyl]-2-(3-aminocarbonylphenyl)-5-  
trifluoromethyl-2*H*-pyrazole-3-carboxamide.

4.2 Analogously to Example 3, reaction of  
2-(3-aminocarbonylphenyl)-5-trifluoromethyl-2*H*-pyrazole-3-  
carboxylic acid with  
30 1-(4-amino-2-chlorophenyl)pyrrolidine-2-thione gives the com-  
pound  
N-[3-chloro-4-(2-thioxopyrrolidin-1-yl)phenyl]-2-(3-amino-  
carbonylphenyl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxamide ("BB"),  
ESI 508.

4.3 Analogously to Example 3, reaction of  
2-(3-aminocarbonylphenyl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxylic acid with  
1-(4-aminophenyl)-2-iminopyrrolidine gives the compound  
N-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-(3-aminocarbonylphenyl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxamide, ESI 457.

### **Example 5**

Analogously to Example 3, reaction of  
2-(3-aminocarbonylphenyl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxylic acid with  
1-(4-aminophenyl)pyrrolidin-2-one *O*-methyl oxime,  
1-(4-amino-2-methylphenyl)pyrrolidin-2-one *O*-methyl oxime,  
gives the compounds  
N-[4-(2-methoxyiminopyrrolidin-1-yl)phenyl]-2-(3-aminocarbonylphenyl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxamide, ESI 487;  
N-[3-methyl-4-(2-methoxyiminopyrrolidin-1-yl)phenyl]-2-(3-aminocarbonylphenyl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxamide, ESI 515.

### **Example 6**

6.1 Analogously to Example 3, reaction of  
2-(3-BOC-aminobenzo[d]isoxazol-5-yl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxylic acid with  
4-(2-iminopyrrolidin-1-yl)phenylamine,  
4-(2-imino-5-methyl-3*H*-1,3,4-thiadiazol-3-yl)phenylamine,  
3-bromo-4-(2-imino-5-methyl-3*H*-1,3,4-thiadiazol-3-yl)phenylamine,

4-(1,5-dimethyl-3-imino-1,2-dihydropyrazol-2-yl)phenylamine,

and subsequent removal of BOC gives the compounds

- 5 N-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-(3-aminobenzo[d]isoxazol-5-yl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxamide,  
N-[4-(2-imino-5-methyl-3*H*-1,3,4-thiadiazol-3-yl)phenyl]-2-(3-amino-  
benzo[d]isoxazol-5-yl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxamide,  
10 N-[3-bromo-4-(2-imino-5-methyl-3*H*-1,3,4-thiadiazol-3-yl)phenyl]-2-(3-aminobenzo[d]isoxazol-5-yl)-5-trifluoromethyl-2*H*-pyrazole-3-carbox-  
amide,  
N-[4-(1,5-dimethyl-3-imino-1,2-dihydropyrazol-2-yl)phenyl]-2-(3-  
15 aminobenzo[d]isoxazol-5-yl)-5-trifluoromethyl-2*H*-pyrazole-3-carbox-  
amide.

- 6.2 Analogously to Example 3, reaction of  
20 2-(3-BOC-aminobenzo[d]isoxazol-5-yl)-5-trifluoromethyl-2*H*-pyra-  
zole-3-carboxylic acid with  
1-(4-aminophenyl)pyrrolidine-2-thione and subsequent removal of  
BOC gives the compound  
N-[4-(2-thioxopyrrolidin-1-yl)phenyl]-2-(3-aminobenzo[d]isoxazol-5-  
25 yl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxamide.

- 6.3 Analogously to Example 3, reaction of  
2-(3-aminocarbonylphenyl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxylic  
30 acid with  
3-bromo-4-(2-imino-5-methyl-3*H*-1,3,4-thiadiazol-3-yl)phenylamine,  
4-(2-imino-5-methyl-3*H*-1,3,4-thiadiazol-3-yl)phenylamine,  
4-(2-iminoimidazolidin-1-yl)phenylamine,  
4-(2-iminoimidazolidin-1-yl)-3-methylphenylamine,  
35 4-(2-cyanoiminoimidazolidin-1-yl)phenylamine,

4-(2-cyanoimino-3-methylimidazolidin-1-yl)phenylamine,  
4-(2-imino-5-ethyl-3*H*-1,3,4-thiadiazol-3-yl)phenylamine,  
4-(2-imino-5-aminocarbonyl-3*H*-1,3,4-thiadiazol-3-yl)phenylamine,  
4-(2-imino-5-ethoxycarbonyl-3*H*-1,3,4-thiadiazol-3-yl)phenylamine,

5

gives the compounds

N-[3-bromo-4-(2-imino-5-methyl-3*H*-1,3,4-thiadiazol-3-yl)phenyl]-2-(3-aminocarbonylphenyl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxamide,  
ESI 567, 568;

10

N-[4-(2-imino-5-methyl-3*H*-1,3,4-thiadiazol-3-yl)phenyl]-2-(3-aminocarbonylphenyl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxamide, ESI 488;

15

N-[4-(2-iminoimidazolidin-1-yl)phenyl]-2-(3-aminocarbonylphenyl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxamide;

N-[4-(2-iminoimidazolidin-1-yl)-3-methylphenyl]-2-(3-aminocarbonylphenyl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxamide;

20

N-[4-(2-cyanoiminoimidazolidin-1-yl)phenyl]-2-(3-aminocarbonylphenyl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxamide ("CA");

N-[4-(2-cyanoimino-3-methylimidazolidin-1-yl)phenyl]-2-(3-aminocarbonylphenyl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxamide;

25

N-[4-(2-imino-5-ethyl-3*H*-1,3,4-thiadiazol-3-yl)phenyl]-2-(3-aminocarbonylphenyl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxamide ("CB"),

N-[4-(2-imino-5-aminocarbonyl-3*H*-1,3,4-thiadiazol-3-yl)phenyl]-2-(3-aminocarbonylphenyl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxamide,

N-[4-(2-imino-5-ethoxycarbonyl-3*H*-1,3,4-thiadiazol-3-yl)phenyl]-2-(3-aminocarbonylphenyl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxamide.

30

6.4 Analogously to Example 3, reaction of

5-(3-aminocarbonylphenyl)-2-methylthiazole-4-carboxylic acid with

4-(2-imino-5-ethyl-3*H*-1,3,4-thiadiazol-3-yl)phenylamine

35

gives the compound.

N-[4-(2-imino-5-ethyl-3*H*-1,3,4-thiadiazol-3-yl)phenyl]-5-(3-amino-carbonylphenyl)-2-methylthiazole-4-carboxamide.

5        6.5        Analogously to Example 3, reaction of  
             2-(3-aminocarbonylphenyl)-5-methyl-2*H*-pyrazole-3-carboxylic acid  
             with 4-(2-imino-5-ethyl-3*H*-1,3,4-thiadiazol-3-yl)phenylamine  
             gives the compound

10        N-[4-(2-imino-5-ethyl-3*H*-1,3,4-thiadiazol-3-yl)phenyl]-2-(3-amino-  
             carbonylphenyl)-5-methyl-2*H*-pyrazole-3-carboxamide.

### **Example 7**

15        7.1        Analogously to Example 3, reaction of  
             2-(3-BOC-amino-1*H*-indazol-5-yl)-5-trifluoromethyl-2*H*-pyrazole-3-  
             carboxylic acid with

20        1-(4-aminophenyl)pyrrolidin-2-one O-methyl oxime,  
             1-(4-aminophenyl)pyrrolidine-2-thione

and subsequent removal of BOC gives the compounds

25        N-[4-(2-methoxyiminopyrrolidin-1-yl)phenyl]-2-(3-amino-1*H*-inda-  
             zol-5-yl)-5-trifluoromethyl-2*H*-pyrazole-3-carboxamide,  
             N-[4-(2-thioxopyrrolidin-1-yl)phenyl]-2-(3-amino-1*H*-indazol-5-yl)-5-  
             trifluoromethyl-2*H*-pyrazole-3-carboxamide.

30

### **Example 8**

             Analogously to Example 3, reaction of  
35        2-(3-thiocarbamoylphenyl)-5-trifluoromethyl-2*H*-pyrazole-3-car-  
             boxylic acid with

1-(4-aminophenyl)pyrrolidin-2-one O-methyl oxime gives the compound

N-[4-(2-methoxyiminopyrrolidin-1-yl)phenyl]-2-(3-thiocarbamoylphenyl)-5-trifluoromethyl-2H-pyrazole-3-carboxamide.

5

### **Example 9**

10

*Preparation of N-[4-(2-hydroxyiminopyrrolidin-1-yl)phenyl]-2-(3-aminomethylphenyl)-5-trifluoromethyl-2H-pyrazole-3-carboxamide*

15

Hydroxylammonium hydrochloride and triethylamine are added to a solution of N-[4-(2-methylsulfanylpyrrolidin-1-yl)phenyl]-2-(3-BOC-aminomethylphenyl)-5-trifluoromethyl-2H-pyrazole-3-carboxamide, iodide in ketone-free ethanol, and the mixture is stirred at room temperature.

20

After 20 hours, the mixture is evaporated to dryness, and the residue is stirred into water and filtered off. The crude product is dried, and 20 ml of HCl in ether are added. After 20 hours, the solvent is removed under reduced pressure, and the product is dried by stirring with ether, giving N-[4-(2-hydroxyiminopyrrolidin-1-yl)phenyl]-2-(3-aminomethylphenyl)-5-trifluoromethyl-2H-pyrazole-3-carboxamide.

25

### **Pharmacological data**

30

Affinity to receptors

Table 1

35

Compound No.	FXa-IC <sub>50</sub> [M]	TF/FVIIa-IC <sub>50</sub> [M]
"AA"	9.6 x 10 <sup>-9</sup>	2.3 x 10 <sup>-8</sup>
"BB"	3.3 x 10 <sup>-7</sup>	1.8 x 10 <sup>-7</sup>

- 47 -

"CA"	$2.0 \times 10^{-7}$	$6.9 \times 10^{-8}$
"CB"	$9.2 \times 10^{-8}$	$1.5 \times 10^{-7}$

5

10

15

20

25

30

35

The following examples relate to pharmaceutical preparations:

5

**Example A: Injection vials**

10

A solution of 100 g of an active ingredient of the formula I and 5 g of disodium hydrogenphosphate in 3 l of bidistilled water is adjusted to pH 6.5 using 2N hydrochloric acid, sterile filtered, transferred into injection vials, lyophilised under sterile conditions and sealed under sterile conditions. Each injection vial contains 5 mg of active ingredient.

15

**Example B: Suppositories**

20

A mixture of 20 g of an active ingredient of the formula I with 100 g of soya lecithin and 1400 g of cocoa butter is melted, poured into moulds and allowed to cool. Each suppository contains 20 mg of active ingredient.

25

**Example C: Solution**

A solution is prepared from 1 g of an active ingredient of the formula I, 9.38 g of  $\text{NaH}_2\text{PO}_4 \cdot 2 \text{H}_2\text{O}$ , 28.48 g of  $\text{Na}_2\text{HPO}_4 \cdot 12 \text{H}_2\text{O}$  and 0.1 g of benzalkonium chloride in 940 ml of bidistilled water. The pH is adjusted to 6.8, and the solution is made up to 1 l and sterilised by irradiation. This solution can be used in the form of eye drops.

30

**Example D: Ointment**

35

500 mg of an active ingredient of the formula I are mixed with 99.5 g of Vaseline under aseptic conditions.



**Example E: Tablets**

5

A mixture of 1 kg of active ingredient of the formula I, 4 kg of lactose, 1.2 kg of potato starch, 0.2 kg of talc and 0.1 kg of magnesium stearate is pressed to give tablets in a conventional manner in such a way that each tablet contains 10 mg of active ingredient.

**Example F: Coated tablets**

10

Tablets are pressed analogously to Example E and subsequently coated in a conventional manner with a coating of sucrose, potato starch, talc, tragacanth and dye.

15

**Example G: Capsules**

20

2 kg of active ingredient of the formula I are introduced into hard gelatine capsules in a conventional manner in such a way that each capsule contains 20 mg of the active ingredient.

**Example H: Ampoules**

25

A solution of 1 kg of active ingredient of the formula I in 60 l of bidistilled water is sterile filtered, transferred into ampoules, lyophilised under sterile conditions and sealed under sterile conditions. Each ampoule contains 10 mg of active ingredient.

30

35